

Power Weakly Mixing Infinite Transformations

Sarah L. Day ^{*} Brian R. Grivna [†] Earle P. McCartney [‡]

Cesar E. Silva [§]

March 1, 1998

Abstract

We construct a rank one infinite measure preserving transformation T such that for all sequences of nonzero integers $\{k_1, \dots, k_r\}$, $T^{k_1} \times \dots \times T^{k_r}$ is ergodic.

AMS 1991 subject classification: 28D.

Key words: ergodic index, weak mixing, infinite measure transformation

^{*}Emory University, Atlanta, GA 30332

[†]St. Olaf College, Northfield, MN 55057

[‡]Williams College, Williamstown, MA 01267

[§]Williams College, Williamstown, MA 01267, csilva@williams.edu

1 Introduction

It is well known that for the case of finite measure preserving transformations, if T is weakly mixing then $T^{k_1} \times \dots \times T^{k_r}$ is ergodic for any sequence of nonzero integers $\{k_1, \dots, k_r\}$. Kakutani and Parry proved in [KP] that there exist infinite (measure preserving) transformations such that $T \times \dots \times T$ (r terms) is ergodic but $T \times \dots \times T$ ($r+1$ terms) is not; in this case the transformation is said to have **ergodic index** r . T is said to have **infinite ergodic index** if it has ergodic index r for all $r > 0$. In [KP], they also constructed infinite Markov shifts of infinite ergodic index. Furthermore, for the case of infinite transformations, it was shown in [ALW] that ergodicity of $T \times T$ implies weak mixing but that there exist infinite weak mixing transformation with $T \times T$ not conservative, hence not ergodic. Later it was shown that T may be weakly mixing with $T \times T$ conservative but still not ergodic, and that there exist rank one infinite transformations of infinite ergodic index [AFS].

In this paper we introduce a condition stronger than infinite ergodic index. Define a transformation T to be **power weakly mixing** if for all finite sequences of nonzero integers $\{k_1, \dots, k_r\}$,

$$T^{k_1} \times \dots \times T^{k_r}$$

is ergodic. Clearly, any power weakly mixing transformation has infinite ergodic index. As T is weakly mixing, it follows that for all ergodic finite measure preserving transformations S , $T \times S$ is ergodic; however, there always exists a conservative ergodic infinite measure preserving transformation R such that $T \times R$ is not conservative, hence not ergodic [ALW]. Recently, it has been shown that infinite ergodic index does not imply power weakly mixing [AFS2].

In section 2 we prove some preliminaries on approximation and in section 3 we construct a rank one infinite measure preserving transformation which is power weakly mixing. We refer to [AFS] for terms not defined here.

Acknowledgments. This paper is based on research in the Dynamical Systems group of the 1997 SMALL Undergraduate Summer Research Project at Williams College with Prof. C. Silva as faculty advisor. Support for the project was provided by a National Science Foundation REU Grant and the Bronfman Science Center of Williams College.

2 Approximation Properties

In this section we prove an approximation lemma for transformations defined by cutting and stacking [F]. This idea has been used earlier in e.g. [AFS] to show that a specific transformation has infinite ergodic index. However, here we present it in greater generality that permits other applications such as in [AFS2]. Thus we first describe cutting and stacking constructions [F].

Let X be a finite or infinite interval of real numbers and μ be Lebesgue measure. A **column** consists of a collection of pairwise disjoint intervals in X of the form $B^0, B^1, \dots, B^{h-1}$, where $\mu(B) > 0$ and $h > 0$. The elements of $\mathcal{C}$ are called **levels** and h is the **height** of $\mathcal{C}$. The column $\mathcal{C}$ partially defines a transformation T on levels B^i , $i = 0, \dots, h-2$, by the translation that takes interval B^i to interval B^{i+1} . Thus sometimes we write B^i as $T^i B$.

A **cutting and stacking** construction for a measure preserving transformation $T : X \rightarrow X$ consists of a sequence of columns

$$\mathcal{C}_n = \{B_n^0, B_n^1, \dots, B_n^{h_n-1}\}$$

of height h_n such that:

i) $\mathcal{C}_{n+1}$ is obtained by cutting $\mathcal{C}_n$ into c_n equal-measure subcolumns or **copies**, putting a number of **spacers** (new levels of the same measure as any of the levels in the c_n subcolumns) above each subcolumn, and stacking from left to right (i.e., the top (or top spacer if it exists) of the left subcolumn is sent by translation to the bottom of its right subcolumn). We assume $c_n \geq 2$. In this way $\mathcal{C}_{n+1}$ consists of c_n copies of $\mathcal{C}_n$, possibly separated by spacers.

ii) B_n is a union of elements from $\{B_{n+1}, TB_{n+1}, \dots, T^{h_{n+1}-h_n} B_{n+1}\}$.

iii) $\bigcup_n \mathcal{C}_n$ generates the Borel sets, i.e., for all subsets A in X , $\mu(A) > 0$, and for all $\epsilon > 0$, there exists C , a union of elements from $\mathcal{C}_n$, for some n , such that $\mu(A \triangle C) < \epsilon$.

Suppose $I = T^j B_k$ is in $\mathcal{C}_k$. For any $n > k$, I is the union of some elements in $\mathcal{C}_n = \{B_n, TB_n, \dots, T^{h_n-1} B_n\}$. We call the elements in this union **sublevels** of I .

Given a real number $0 < \epsilon < 1$, and a subset A of X with $\mu(A) > 0$, we say that a subset I of X is $(1 - \epsilon)$ -**full** of A provided

$$\mu(I \cap A) > (1 - \epsilon)\mu(I).$$

A set I in the product space $\Pi_{i=1}^r X$ is a **rectangle** if I can be written as the Cartesian product of levels in some column $\mathcal{C}_k$. We let ν be the product measure μ^r . Rectangles I are defined to be $(1 - \epsilon)$ -full of a set A in a similar way as before.

Lemma 2.1. *Given subsets A and B of $\Pi_{i=1}^r X$, and $\epsilon > 0$, there exist rectangles $I = I_1 \times \dots \times I_r$ and $J = J_1 \times \dots \times J_r$ with $I_1, \dots, I_r, J_1, \dots, J_r$ in a column $\mathcal{C}_k$ such that for all $m = 1, \dots, r$, I_m may be chosen to be either above or below J_m and with I and J $(1 - \epsilon)$ -full of sets A and B respectively.*

Proof. Choose rectangles $I' = I'_1 \times \dots \times I'_r$ and $J' = J'_1 \times \dots \times J'_r$, with I'_m and J'_m in $\mathcal{C}_{k-1}$, such that I' and J' are $(1 - \frac{\epsilon}{c_{k-1}^r})$ -full of A and B respectively. Now look at any two copies of $\mathcal{C}_{k-1}$ in $\mathcal{C}_k$. To have I_m above J_m , let I_m be the top copy of I'_m in $\mathcal{C}_k$ and let J_m be the bottom copy of J'_m in $\mathcal{C}_k$. To have I_m below J_m make an analogous choice. Let $I = I_1 \times \dots \times I_r$ and $J = J_1 \times \dots \times J_r$. One verifies that I and J are $(1 - \epsilon)$ -full of A and B . $\square$

Lemma 2.2. (Double Approximation Lemma) *Suppose A is a subset of the product space $\Pi_{i=1}^r X$ with $\nu(A) > 0$. Let $I = I_1 \times \dots \times I_r$ be a rectangle in $\mathcal{C}_k$ that is $(1 - \epsilon)$ -full of A . For $n > k$, let $P_n = c_k \cdots c_{n-1}$, let V_n index the P_n copies of $\mathcal{C}_k$ in $\mathcal{C}_n$, and let $V = V(n) = V_n \times \dots \times V_n$ (r times). Then for any δ , $0 < \delta < 1$, and for any τ , $0 < \tau < 100(1 - \epsilon)$, there exists an integer N such that for all $n > N$, there is a set V'' of size at least τ percent of V such that for all $v = (v_1, \dots, v_r) \in V''$, I_v is $(1 - \delta)$ -full of A and each I_v is of the form $I_v = I''_1 \times \dots \times I''_r$ where I''_m is a sublevel of I_m in the v_m -copy of $\mathcal{C}_k$.*

Proof. For convenience, let A denote $I \cap A$ and let t denote $\frac{\tau}{100}$. Then $\nu(I \triangle A) < \epsilon \nu(I)$. We have that $V_n = \{1, \dots, P_n\}$ and $V = \{(v_1, \dots, v_r) | v_i \in V_n\}$. Then $I = \cup_{v \in V} I_v$.

Choose $c > \frac{\delta+1}{1-t-\epsilon} > 0$. Next pick $N > k$ sufficiently large so that for any $n \geq N$ there exists V' a subset of V such that $I' = \cup_{v \in V'} I_v$ satisfies

$$\nu(I' \triangle A) < \frac{\delta}{c} \nu(I).$$

Thus,

$$\begin{aligned}
\nu(I' \triangle I) &< \frac{\delta}{c}\nu(I) + \epsilon\nu(I) \\
&= \left(\frac{\delta}{c} + \epsilon\right)\nu(I).
\end{aligned}$$

Now let $V'' = \{v \in V' | \nu(I_v \triangle A) < \delta\nu(I_v)\}$ and set $I'' = \cup_{v \in V''} I_v$, the union of the $(1 - \delta)$ -full I_v subintervals.

Then,

$$\begin{aligned}
\delta\nu(I' \triangle I'') &= \sum_{v \in V' \triangle V''} \delta\nu(I_v) \\
&\leq \sum_{v \in V' \triangle V''} \nu(I_v \triangle A) \\
&\leq \nu(I' \triangle A).
\end{aligned}$$

So,

$$\begin{aligned}
\nu(I'' \triangle I) &\leq \frac{1}{\delta}\nu(I' \triangle A) + \nu(I' \triangle I) \\
&< \frac{1}{c}\nu(I) + \left(\frac{\delta}{c} + \epsilon\right)\nu(I) \\
&< (1 - t)\nu(I).
\end{aligned}$$

Therefore, more than τ percent of the subrectangles contained in I are in I'' and are thus $(1 - \delta)$ -full of A . $\square$

3 A Power Weakly Mixing T

In this section we construct a rank one infinite measure preserving transformation T that is power weakly mixing; then we mention how the proof gives a family of such transformations. We start by defining inductively a sequence of columns $\{\mathcal{C}_n\}$. Let $\mathcal{C}_1$ have base $B_1 = [0, 1)$ and height $h_1 = 1$. Given a column $\mathcal{C}_k$ with base $B_k = [0, \frac{1}{4^k-1})$ and height h_k , $\mathcal{C}_{k+1}$ is formed by cutting $\mathcal{C}_k$ vertically three times so that B_k is cut into the intervals $B_{k,1} = [0, \frac{1}{4^k})$, $B_{k,2} = [\frac{1}{4^k}, \frac{1}{2}(\frac{1}{4^k-1}))$, $B_{k,3} = [\frac{1}{2}(\frac{1}{4^k-1}), \frac{3}{4}(\frac{1}{4^k-1}))$, $B_{k,4} = [\frac{3}{4}(\frac{1}{4^k-1}), \frac{1}{4^k-1})$. We then add a column of spacers h_k high to the top of the subcolumn whose base is $B_{k,2}$. Next we add one spacer to the top of the subcolumn




Figure 1: A $\mathcal{C}_n$ column.

whose base is $B_{k,4}$; this is called the **staircase spacer** of $\mathcal{C}_k$. Then stack from left to right, i.e., the top level on the left is sent to the bottom level on the right by the translation map. The resulting column $\mathcal{C}_{k+1}$ now has base $B_{k+1} = [0, \frac{1}{4^k})$ and height $h_{k+1} = 5h_k + 1$. The union of the columns is $X = [0, \infty)$. This defines a conservative ergodic rank one infinite measure preserving transformation T .

Any column $\mathcal{C}_n = \{B_n, \dots, T^{h_n-1}B_n\}$ has four subcolumns $\mathcal{C}_{n,i} = \{B_{n,i}, \dots, T^{h_n-1}B_{n,i}\}$ for $i = 1, \dots, 4$. Given a level L in $\mathcal{C}_n$ and an integer $\ell > 0$, we will be interested in studying $T^{\ell h_n}L$ (a translation of L through $\mathcal{C}_n$ ℓ times). To simplify our estimates, we will only be concerned with the part of $T^{\ell h_n}L$ that is in $\mathcal{C}_{n,1}$; this will consist of a sequence of sublevels that we call an **L -crescent** (refer to figure 1).

Theorem 3.1. *For any sequence of nonzero integers $\{k_1, \dots, k_r\}$, the transformation $T^{k_1} \times \dots \times T^{k_r}$ is ergodic.*

Proof. Let A and B be in $\Pi_{i=1}^r X$ with $\nu(A) > 0$ and $\nu(B) > 0$. Find rectangles $I = I_1 \times \dots \times I_r$ and $J = J_1 \times \dots \times J_r$ such that

$$\nu(A \cap I) > \frac{3}{4}\nu(I),$$

$$\nu(B \cap J) > \frac{3}{4}\nu(J),$$

and $I_m, J_m, m = 1, \dots, r$ are all in the same column $\mathcal{C}_k$, and I_m is above J_m if k_m is positive, and I_m is below J_m if k_m is negative.

Suppose L is a level in $\mathcal{C}_n$ for any $n \geq k$. Translating L by some multiple ℓh_n of the height of the column results in crescents. It suffices to consider a worst case lower bound to various intersections, thus we will only consider crescents in the leftmost subcolumn $\mathcal{C}_{n,1}$. The minimum size of the top of any such crescent is at least $\frac{1}{8^\ell}\mu(L)$ (given the nature of our construction, the size of the crescent may not decrease after each step, but to simplify our calculations we use a conservative estimate). In addition, each crescent has moved through the staircase spacers ℓ times. The maximum total number of staircase spacers that any given crescent has moved through is $s_\ell = \sum_{i=1}^\ell i$. Therefore, any level J more than s_ℓ below I contains some pieces of the crescent from I . Furthermore, the minimum amount of the crescent from I that intersects J has measure at least $\frac{1}{8^{\ell+d}}\mu(J)$, where d is the distance J is below I .

To account for the fact that after each pass through a staircase spacer the crescent is moved down by one, we note that translating I by $\ell h_n + c$, with $c > s_\ell$, ensures that any level J below (or at) I contains pieces of the I -crescent having total measure at least $\frac{1}{8^{\ell+d+c}}\mu(J)$, where d is as above.

For each k_i , let $s_{k_i} = \sum_{j=1}^{k_i} j$, let d_i be the distance between I_i and J_i for all i , and put $K = \max\{k_i\}$, $S = \max\{s_{k_i}\}$, and $d = \max\{d_i\}$. Choose δ so that

$$0 < \delta < \left(\frac{1}{8^{K+d+KS}}\right)^r.$$

By the Double Approximation Lemma find $I' = I'_1 \times \dots \times I'_r$ and $J' = J'_1 \times \dots \times J'_r$ such that I' and J' are $(1 - \delta)$ -full of A and B respectively, $I'_1, \dots, I'_r, J'_1, \dots, J'_r$ are all in some column $\mathcal{C}_n$, and for each i , I'_i and J'_i are in the same $\mathcal{C}_k$ -copy in $\mathcal{C}_n$, and I_i is more than SK levels below the top of $\mathcal{C}_n$.

Let $H = h_n + S$. Then for all positive k_i ,

$$\mu(T^{k_i H} I'_i \cap J'_i) \geq \frac{1}{8^{k_i+d_i+k_i s_i}} \mu(I'_i) \geq \frac{1}{8^{K+d+KS}} \mu(I'_i).$$

For all negative k_i ,

$$\mu(T^{k_i H} I'_i \cap J'_i) = \mu(I'_i \cap T^{|k_i| H} J'_i) \geq \frac{1}{8^{k_i+d_i+k_i s_i}} \mu(J'_i) \geq \frac{1}{8^{K+d+KS}} \mu(J'_i).$$

Therefore,

$$\nu[(T^{k_1} \times \dots \times T^{k_r})^H I' \cap J'] \geq (\frac{1}{8^{K+d+KS}})^r \nu(I').$$

Thus,

$$\begin{aligned} \nu[(T^{k_1} \times \dots \times T^{k_r})^H A \cap B] &\geq \nu[(T^{k_1} \times \dots \times T^{k_r})^H I' \cap J'] - \delta \nu(I') \\ &\geq (\frac{k}{8^{K+d+KS}})^r \nu(I') - \delta \nu(I') > 0. \end{aligned}$$

Therefore $T^{k_1} \times \dots \times T^{k_r}$ is ergodic. □

Remark. 1. The same proof will apply to a transformation where at the k^{th} stage column C_k is cut into $c > 1$ equally-spaced subcolumns $\mathcal{C}_{k,1}, \dots, \mathcal{C}_{k,c}$, a single (stair-case) spacer is put on top of column $\mathcal{C}_{k,c}$ and a stack of h_k spacers is put on top of any of the middle subcolumns.

2. There exists a rank one infinite measure preserving transformation S such that S has infinite ergodic index but $S \times S^2$ is not conservative, hence S is not power weakly mixing [AFS2].

References

- [ALW] J. Aaronson, M. Lin, and B. Weiss, *Mixing properties of Markov operators and ergodic transformations, and ergodicity of Cartesian products*, Israel J. Math. 33, 1979, 198-224.
- [AFS] T. Adams, N. Friedman, and C.E. Silva, *Rank-one weak mixing for nonsingular transformations*, Israel J. Math. 102 (1997), 269-281.
- [AFS2] T. Adams, N. Friedman, and C.E. Silva, *Rank-one weak mixing for nonsingular transformations II*, preprint.
- [F] N.A. Friedman, *Introduction to Ergodic Theory*, Van Nostrand, 1970.
- [KP] S. Kakutani and W. Parry, *Infinite measure preserving transformations with “mixing”*, Bull. Amer. Math. Soc. 69, 1963, 752-756.